

Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Delvaeye M, Noris M, De Vriese A, et al. Thrombomodulin mutations in atypical hemolytic–uremic syndrome. *N Engl J Med* 2009;361:345-57.

Supplementary Methods

Patients and controls

All patients with aHUS with available stored DNA, and who had been registered consecutively with the International Registry of Recurrent and Familial HUS/TTP were recruited into this study. The controls were not individually matched. We used two different cohorts of controls. The first cohort comprised 150 healthy controls from the International Registry of Recurrent and Familial HUS/TTP²⁷. They were selected for having a median age and female/male ratio and geographic origin (125 from Europe; 20 from North America, white; 5 from Africa) comparable to patients. The geographic origin of patients was as follows: 58% from Italy, 21% from other European countries, 15% from North America, 2% from South America, 2% from Africa, and 1.3% from Middle East countries. To further confirm that the first cohort was representative of a wider population, we studied an additional 230 available DNAs from a second cohort of healthy white caucasians from Europe, relatives of patients with a range of diseases, with a median age older than the first cohort, who had voluntarily donated blood into a databank for research purposes.

aHUS was diagnosed in patients who had one or more episodes of microangiopathic hemolytic anemia and thrombocytopenia (hematocrit <30%, hemoglobin <10 g/dl, serum lactate dehydrogenase >460 U/L, undetectable haptoglobin, fragmented erythrocytes in the blood smear, and platelet count <150,000/ μ l), associated with acute renal failure. The latter was defined according to the RIFLE classification²⁸, as an increase in serum creatinine x 3.0 or a decrease in glomerular filtration rate by 75%, or serum creatinine

levels ≥ 4 mg/dL with an acute increase of more than 0.5 mg/dL; urine output < 0.3 ml/kg/h for 24 hours or anuria for 12 hours.

Patients with Stx-HUS (defined by the presence of free fecal Stx and/or by positive colony sweeps for Stx production in MacConkey agar culture of stools, by Vero cell assay, and/or serum antibodies measured by ELISA to Stx and/or to lipopolysaccharide of the five major Stx-producing serogroups, O157, O26, O103, O111 and O154⁵⁸) were excluded.

Of the 152 patients, 18 had a familial history of the disease, of whom 3 were from one pedigree and 2 from another one. The others were unrelated, including F635 and F163. For the latter two patients in whom thrombomodulin mutations were found, all available relatives were thereafter recruited and screened. All the other patients had the sporadic form of aHUS and were all unrelated. The 380 controls were unrelated to each other and unrelated to patients.

DNA sequencing and genotyping of SNPs

Genomic DNA was extracted from peripheral blood leukocytes. The entire coding sequence of *THBD*, including the 5' and 3' untranslated regions and approximately 50 bp of 5' and 3' flanking DNA, was directly sequenced on both strands, using 5 pairs of oligonucleotide primers. Sequences that indicated the presence of SNPs were visually confirmed on both strands, and repeated with fresh DNA samples. Further confirmation of SNPs was achieved using the Sequenom MALDI TOF MassArray system³¹. Thus, for SNP analysis, genotyping assays (primers for PCR amplification and the extension probe) were designed using the Sequenom MassARRAY Assay

Design 3.0 software, applying default parameters. Multiplexed PCR was performed with 0.2 units of Taq polymerase and 20 ng of genomic DNA. Thermocycling was at 95°C for 15 min followed by 45 cycles of 95°C for 20s, 56°C for 30s and 72°C for 1 min. Unincorporated dNTPs were deactivated using shrimp alkaline phosphatase, and primer extension was carried out using extension primers, ddNTPs and Thermosequenase DNA polymerase. Reactions were cycled at 94°C for 2 min, followed by 44 cycles of 94°C for 5s, 52°C for 5s and 80°C for 5s. After the addition of a cation exchange resin to remove residual salt from the reactions, the purified primer extension reaction was loaded onto a matrix pad (3-hydroxypicolinic acid) of a SpectroCHIP (Sequenom) and analyzed using a matrix-assisted laser desorption/ionization–time of flight (MALDI-TOF) mass spectrometer.

DNA preparation and cell transfections

Genomic DNA representing the entire coding region of human thrombomodulin was generated by PCR and subcloned into the *EcoR1* sites of the expression vector pcDNA3.1 (Invitrogen). Mutations were introduced with the Quikchange XL site-directed mutagenesis kit (Stratagene, Amsterdam, the Netherlands). Transfections into cultured mammalian cells were performed with Fugene HD (Roche, Brussels, Belgium) in 24-well plates with 0.2 ug DNA per well. Selection was achieved with neomycin (G418, Roche, Belgium) and cells were maintained in DMEM/F12 (CHO-K1) or DMEM (HEK293), supplemented with 10% fetal calf serum (Invitrogen, Merelbeke, Belgium).

Proteins and antibodies

SDS-PAGE and Western immunoblotting was performed by standard techniques. Complement reagents were from Complement Technology, Inc. (Tyler, Texas, USA). Purified proteins were biotinylated with 50x molar excess D-Biotinoyl-e-amidocaproic acid-N-hydroxysuccinimide ester (Boehringer Mannheim, Brussels, Belgium) in PBS and dialyzed to remove free biotin. Murine monoclonal antibodies (CTM1009) were generated against chondroitin sulfate free human extramembranous thrombomodulin (RSV TM)⁵⁹ by standard procedures, purified using Med Hypercel (VWR, USA). The following primary antibodies were used: goat anti-human C3, and goat anti-human CFH. Sheep anti-human C4bBP was from Abcam (Cambridge, UK).

Co-precipitation assays

Proteins that interact with recombinant thrombomodulin (RSV TM)⁵⁹, were co-precipitated with sepharose-bound human diisopropylfluorophosphate (DFP)-inactivated thrombin (Enzyme Research Labs, IN) after incubation with RSVTM.

Detection of C3b proteolytic fragments

Recombinant thrombomodulin was incubated with different combinations of C3b 0.5 µg, CFH 100 ng, C4bBP 100 ng, and CFI 100 ng in a volume of 150 µl of PBS at 37°C for 90'. After reactions and separation by SDS-PAGE, C3b proteolytic fragments were analysed by Western blot and quantified by densitometry. The same experiments were performed on the surface of HEK293 cells transfected with pcDNA3.1 (empty vector) or vector expressing thrombomodulin (full-length human thrombomodulin). Equal expression of

thrombomodulin in transfected cells was confirmed by Western blot and by ELISA.

Thrombin-mediated TAFI activation

HEK293 cells were incubated at 37°C for 20 mins with 3 ug of TAFI in 150 ul of PBS with 1 mM CaCl₂ and 4 U/ml of thrombin. Reactions were stopped with hirudin and the proteins were separated by SDS-PAGE followed by Western immunoblotting to detect TAFIa.

Complement binding assays

Binding of complement proteins to immobilized thrombomodulin was performed with a 2-step procedure. The concentration of thrombomodulin in cell lysates was first quantified using a commercial sandwich ELISA kit (Abcam, Cambridge, UK), but replacing the second anti-thrombomodulin antibody with biotinylated CTM1009. Lysates with 10 ug/ml thrombomodulin were then co-incubated with 3 ug/ml biotinylated C3b or CFH for exactly 3 hr at room temperature in the wells of the commercial thrombomodulin ELISA plates, which are precoated with anti-thrombomodulin antibodies. After washes, specific binding was quantified with a streptavidin-secondary antibody diluted in the strep-HRP diluent and TNB substrate. Reactions were stopped with H₂SO₄. Absorbance was measured at 495 nm with an ELISA plate reader (Beun De Ronde, Serlabo, Belgium). Non-specific signal, determined from parallel experiments with lysates of cells transfected with empty vector (no thrombomodulin), was subtracted from the result. In all cases, co-incubation of biotinylated protein with a molar excess of the

corresponding non-biotinylated protein abrogated the specific signal (not shown). Assays were performed three times in triplicate.

Initiation of complement activation and flow cytometry

Initiation of the complement pathway on CHO-K1 cells was performed as described^{32, 33}. CHO-K1 cells stably expressing equal amounts of cell-surface wild-type or mutant forms of thrombomodulin were trypsinized with 0.05% trypsin in PBS, washed in 1% FCS/PBS, and sensitized with anti-CHO antibodies (Cygnus, Southpark, NC, USA) for 30' at 4°C. After washing in gelatin veronal buffer (GVB++) (Complement Technologies Inc., Tyler, Texas, USA), cells were incubated at 37°C for 90' in 10% C6-deficient serum (which allows complement activation to proceed without inducing cell lysis) diluted in GVB ++ buffer, washed twice with 1% FCS/PBS, and incubated at 4°C for 30 minutes with 20 ug/ml of murine monoclonal anti-thrombomodulin antibodies (CTM1009), FITC-conjugated murine anti-human C3c antibodies (11.3 ug/ml, Dakocytomation, Heverlee, Belgium, Heverlee) (identifies C3b and iC3b), or murine anti-human iC3b antibodies (10 ug/ml, Quidel, Osteomedical, the Netherlands, Nijkerk). Species and isotype matched immunoglobulins were used as negative controls. After incubation with an appropriate secondary antibody, cells were washed with 1% FCS/PBS, and fixed with 0.5% paraformaldehyde prior to analysis with the FACScalibur (BD Biosciences, Erembodegem, Belgium). The fold-change in % inactivation of C3b (mean fluorescence intensity of iC3b/C3c) was calculated for wild-type and mutant thrombomodulin-expressing cells, relative to that in non-thrombomodulin expressing cells (i.e. empty vector). Reported results reflect the mean \pm standard error of the mean (SEM) of 3 independent experiments.

Statistical analyses

Data analyses were performed with Graphpad Prism 4 (San Diego, CA, USA). Results are presented as mean \pm standard error of the mean (SEM). An unpaired t-test was used to identify differences between groups. Allele frequencies of *THBD* gene variants in patients and controls were compared by the chi-square test. All reported P-values are 2-sided and not adjusted for multiple testing.

Author statements

EM Conway, M Delvaeye, M Noris, G Remuzzi, D Lambrechts, CT Esmon and NL Esmon designed the studies and contributed to all versions of the manuscript. EM Conway and M Noris wrote the major portion of the first draft. M Delvaeye, B Claes, J Del-Favero, S Plaisance and C Zoja performed experiments, gathered and analyzed data. EM Conway, M Noris and D Lambrechts validated data. All authors agreed with submission of the manuscript.

SNP	Location relative to ATG start site of <i>THBD</i>	Genotypes and Alleles			p-value
			Controls n=648	HUS n=152	
			Number (%)	Number (%)	
rs1040585	9.8 kb 5'	AA	5 (1)	0 (0)	0.23
		AC	62 (11)	9 (7)	
		CC	483 (88)	115 (93)	
		A	72 (7)	9 (4)	0.08
		C	1028 (93)	239 (96)	
rs2424505	5.75 kb 5'	AA	9 (2)	0 (0)	0.33
		AG	94 (16)	19 (15)	
		GG	479 (82)	109 (85)	
		A	112 (10)	19 (7)	0.27
		G	1052 (90)	237 (93)	
rs6076016	4.3 kb 5'	AA	66 (11)	11 (8)	0.21
		AT	265 (45)	56 (40)	
		TT	261 (44)	72 (52)	
		A	397 (34)	78 (28)	0.08
		T	787 (66)	200 (72)	
rs1042580	2.7 kb 3' (in 3'UTR)	AA	217 (36)	52 (34)	0.6
		AG	285 (47)	78 (52)	
		GG	99 (16)	21 (14)	
		A	719 (60)	182 (60)	0.89
		G	483 (40)	120 (40)	
rs3176123	2.9 kb 3' (in 3'UTR)	AA	396 (69)	108 (76)	0.22
		AC	164 (28)	31 (22)	
		CC	17 (3)	3 (2)	
		A	956 (83)	247 (87)	0.09
		C	198 (17)	37 (13)	
rs1962	3.8 kb 3' (in 3'UTR)	CC	35 (6)	9 (5)	0.82
		CT	202 (34)	51 (34)	
		TT	355 (60)	92 (61)	
		C	272 (23)	69 (23)	0.92
		T	912 (77)	235 (77)	
rs1042579	exon (A473V)	CC	299 (79)	118 (78)	0.43
		CT	73 (19)	33 (22)	
		TT	8 (2)	1 (1)	
		C	671 (88)	269 (88)	0.93
		T	89 (12)	35 (12)	

Supplementary Table 1. Genotype and allele frequencies of common SNPs in *THBD* gene in patients with aHUS and in healthy subjects. SNPs in the *THBD* gene were selected, based on the HAPMAP, using the SNP tagged approach (SNPs with frequency > 0.05 and $R^2 \geq 0.8$) and genotyped using the Sequenom Massarray technology. P-values were calculated using standard chi squared testing, with 1 and 2 degrees of freedom for alleles and genotypes, respectively. Genotype frequencies of SNPs in the controls and cases followed Hardy-Weinberg equilibrium. We also found rare flanking SNPs, 2 controls with G->C at base position -57, 1 control with C->T at base position -11, 2 controls with G->T at base position 2049, and 6 controls and 3 patients with A->G at base position 2079. All of these were heterozygous.

Supplementary Appendix 1

Case Reports

F635, pedigree #185. The proband is a 24 year-old man who had four episodes of HUS between 1 and 3 years of age, that led to end stage renal disease. The patient is now stable and on chronic hemodialysis. None of the episodes was preceded with a diarrhea prodrome. He had five brothers and three sisters. Three of the brothers (including twins) had HUS in early childhood and died during acute episodes of the disease. A younger sister, now 15 years old, also had an episode of HUS during infancy, but she recovered renal function, although she has persistent proteinuria. None of the proband's affected siblings had diarrhea prodromes. There are two older brothers and two older sisters who have never had HUS and they are alive and well. There is no known consanguinity between the parents, who are also healthy.

The proband was screened for complement genes (*CFH*, *MCP*, *CFI*, *C3* and *CFB*) and no mutations were found. Serum C3 was slightly reduced (82 mg/dl, normal values 90-180 mg/dl) and C4 concentration (30 mg/dl, normal values 10-40 mg/dl) was normal. Serum CFH levels were normal (515.6 mg/l, normal values 350-750 mg/l).

F163, pedigree #008. The proband is an 8 year-old boy and is the youngest of three brothers. The oldest died at the age of 8 months of HUS. The second sibling is a 10 year old boy. At present, this brother and the parents are healthy. The proband had three episodes of HUS, between 6 months and 2

years of age. All episodes were triggered by upper respiratory tract infections and characterized by acute renal failure, hypertension, hematuria and proteinuria. The boy was treated with plasmapheresis, blood transfusions and hemodialysis, always obtaining complete remissions. The proband, his healthy brother and the parents were screened for complement genes (*CFH*, *MCP*, *CFI*, *C3* and *CFB*) and no mutations were found. Serum C3 concentration was normal on one occasion (126 mg/dl) and slightly reduced (85 mg/dl) on another. Serum C4 concentrations were normal on both occasions (30 and 28 mg/dl). CFH levels (492 mg/l) were also normal. The activity of the vWF-cleaving protease, ADAMTS13, was normal (activity 110%, normal values 50-150%).

S884. The proband is a 4 year-old boy who developed HUS at the age of 15 months, after a febrile viral illness. The episode was characterized by anemia, thrombocytopenia, increased LDH, with mildly increased serum creatinine (0.6 mg/dl) and proteinuria. Serum C3 (68 mg/dl) was low, while C4 (11 mg/dl) was in the normal range. The boy was treated with plasma infusions and blood transfusions, obtaining a complete remission. He was screened for complement genes (*CFH*, *MCP*, *CFI*, *C3* and *CFB*). On our initial screen, an apparent heterozygous mutation in exon 23 of CFH (C→T at base 3572 causing a S1191L mutation) was detected. However, additional sequencing with different primers confirmed that no mutation existed in CFH. The original error was due to regions of high homology between CFH and CFHL genes. CFH levels (417 mg/L) were normal. No mutations were found in MCP and

CFI. ADAMTS13 activity was normal (81%). There is no relevant family history. The parents are from Yemen.

S511. The proband is a 10 year-old girl who developed HUS at the age of 3 years. She had two consecutive episodes of HUS with severe anemia and thrombocytopenia, and renal impairment (serum creatinine 0.7-1.4 mg/dl), both associated with low C3 levels (40-47 mg/dl). She responded well to plasmapheresis recovering normal renal function, but with persistent mild hypertension. Neither episode was associated with diarrhea or pneumococcal infections. The girl was screened for complement genes (*CFH*, *MCP*, *CFI*, *C3* and *CFB*) and no mutations were found. There is no relevant clinical history in the family.

S015. The proband is a 15 year-old boy who presented with fulminant HUS at the age of 6 months, becoming dialysis-dependent. At the age of 6 years he received a cadaveric kidney transplant, but HUS recurred in the allograft three days after the transplant, with loss of the kidney. Since that time the patient remains on dialysis. Serum C3 levels were normal (92 mg/dl), whereas C4 levels were slightly increased (49 mg/dl). CFH levels (418 mg/l) and ADAMTS13 activity (99%) were normal. The mother died a few years ago due to complications of idiopathic pulmonary fibrosis associated with fulminant pulmonary hypertension. The proband and his healthy father were screened for complement genes (*CFH*, *MCP*, *CFI*, *C3* and *CFB*) and no mutations were found.

S665. The proband is a 23 year old man from Italy who had an episode of HUS at the age of 6 years. He was treated with plasma exchange and recovered. Since then, he has no further episodes. At the last follow-up in 2004, his renal function was borderline normal (serum creatinine 1.24 mg/dl). The patient was screened for *CFH*, *MCP*, *CFI*, *C3* and *CFB*, and no mutations were found. Plasma ADAMTS13 activity are normal (78%). C3 (94 mg/dl), C4 (14 mg/dl) and CFH (515 mg/l) levels are normal.

S924. The patient is a 19 year old woman from the USA, who developed aHUS at the age of 15 years, a few days after receiving a cadaveric kidney transplant. She had severe microangiopathic hemolytic anemia and thrombocytopenia. A biopsy of the graft revealed typical features of thrombotic microangiopathy. Graft function rapidly deteriorated and she developed chronic renal failure. The patient was adopted. The adoptive parents did not have any information on the cause of the primary kidney disease, or whether there was a family history of aHUS. The patient was screened for *CFH*, *MCP*, *CFI*, *C3* and *CFB*, and no mutations were detected. C3 (106 mg/dl) and C4 (30 mg/dl) levels are normal. CFH (949 mg/l) levels are slightly elevated.

Supplementary Appendix 2

International Registry of Recurrent and Familial HUS/ TTP

Coordinators

G. Remuzzi, MD, P. Ruggenenti, MD (Clinical Research Center for Rare Diseases "Aldo e Cele Daccò", Ranica, Bergamo, and Division of Nephrology and Dialysis, "Ospedali Riuniti" Azienda Ospedaliera, Bergamo), **M. Noris, Chem. Pharm. D.** (Clinical Research Center for Rare Diseases "Aldo e Cele Daccò", Ranica, Bergamo).

Investigators - Italy

M. Garozzo, MD (Division of Nephrology and Dialysis, "S. Marta e S. Venera" Hospital, Acireale, Catania); **F. Casucci, MD, F. Cazzato, MD** (Division of Nephrology, "Miulli" Hospital, Acquaviva delle Fonti, Bari); **I.M. Ratsch, MD** (Pediatric Clinic, "G. Salesi" Hospital, Ancona); **S. Alloatti, MD, V. Pellu, MD** (Division of Nephrology and Dialysis, Ospedale Regionale, Aosta); **G. Cesano, MD** (Division of Nephrology, Ospedale Civile, Asti); **G. Claudiani, MD** (Division of Hematology, "S. Liberatore" Hospital, Atri, Teramo); **W. De Simone, MD** (Division of Nephrology and Dialysis, "S. Giuseppe Moscati" Hospital, Avellino); **P. Dattolo, MD, F. Pizzarelli, MD** (Division of Nephrology and Dialysis, "S. M. Annunziata" Hospital, Bagno a Ripoli, Firenze); **R. Bellantuono, MD, T. De Palo, MD** (Division of Nephrology and Dialysis, "Giovanni XXIII" Pediatric Hospital, Bari); **M. Schiavoni, MD** (Assistenza Emofilici e Coagulopatici, Ospedale Policlinico Consorziale, Bari); **M.R. Caruso, MD, E. Gotti, MD, S. Rota, MD, A. Schieppati, MD** (Division of Nephrology and Dialysis, "Ospedali Riuniti" Azienda Ospedaliera, Bergamo); **T. Barbui, MD, A. Falanga, MD** (Division of Hematology, "Ospedali Riuniti" Azienda Ospedaliera, Bergamo); **P. Cornelli, MD, G. Torre, MD** (Pediatric Department, "Ospedali Riuniti" Azienda Ospedaliera, Bergamo); **R. Fumagalli, MD, I. Pelliccioli, MD** (Pediatric Acute Care, "Ospedali Riuniti" Azienda Ospedaliera, Bergamo); **I.M. Berto, MD** (Division of Nephrology, "Ospedale degli Infermi", Biella); **P. Riegler, MD** (Division of Nephrology and Hematology Service, Hospital of Bolzano, Bolzano); **J. Mahlknecht, MD, M. Neunhauserer, MD** (Division of Pediatrics, Hospital of Brunico, Bolzano); **E. Ragazzoni, MD** (Division of Nephrology and Dialysis, Hospital of Borgomanero, Novara); **A.M. Acquarolo, MD** (Il Rianimazione "Spedali Civili, Azienda Ospedaliera", Brescia); **O. Carli, MD, G. Gregorini, MD, S. Bove, MD** (Division of Nephrology and Dialysis, "Spedali Civili, Azienda Ospedaliera", Brescia); **G. Rossi, MD** (Division of Hematology, "Spedali Civili, Azienda Ospedaliera", Brescia); **A. Giangrande, MD** (Division of Nephrology and Dialysis, Ospedale di Circolo, Busto Arsizio, Varese); **P. Founduli, MD** (Pediatric Division, "G. Brotzu" Hospital, Cagliari); **A. Bonadonna, MD, C. Gardin, MD** (Division of Nephrology and Dialysis, Presidio Ospedaliero di Camposampiero, Padova); **A. Mazzotta, MD** (Division of Nephrology and Dialysis, "S. Spirito" Hospital, Casale Monferrato, Alessandria); **G. Delfino, MD, E. Favaro, MD, R. Lazzarin, MD** (Division of Nephrology and Dialysis, "S. Giacomo" Hospital, Castelfranco Veneto, Treviso); **A. Granata, MD** (Division of Nephrology and Dialysis, "Vittorio Emanuele II - Ferrarotto - S. Bambino" Hospitals, Catania); **P. Pignataro, MD** (Division of Medicine, "Vittorio Emanuele II - Ferrarotto - S. Bambino" Hospitals, Catania); **S. Li Volti, MD** (Pediatric Department, Policlinico Hospital, Catania); **M. Rocchietti, MD** (Division of Nephrology and Dialysis, "Ospedale Civile, Ciriè, Torino); **C. Castellino, MD** (Division of Hematology, "Azienda Ospedaliera S. Croce e Carle", Cuneo); **L. Calacoci, MD** (Division of Immunohematology, "S. Giovanni

di Dio" Hospital, Firenze); **C. Grimaldi, MD** (Division of Internal Medicine and Nephrology, "S. Giovanni di Dio" Hospital, Firenze); **I. Pela, MD, D. Seracini, MD** (Division of Nephrology, "A. Meyer" Hospital, Firenze); **R. Piperno, MD, M. Salvadori, MD** (Division of Nephrology and Dialysis, "Careggi" Hospital, Firenze); **E. Capussela, MD** (Division of Hematology, "Ospedali Riuniti" di Foggia, Foggia); **D.A. Procaccini, MD** (Division of Nephrology and Dialysis, "Ospedali Riuniti" di Foggia); **G.C. Barbano, MD, A. Canepa, MD, M.L. Degl'Innocenti, MD, G. Piaggio, MD, A. Trivelli, MD** (Division of Nephrology, "G. Gaslini" Pediatric Institute, Genova); **I. Fontana, MD** (Transplant Center, "S. Martino" Hospital, Genova); **D. Rolla, MD** (Division of Nephrology, Dialysis and Transplantation, "S. Martino" Hospital, Genova); **S. D'Ardia, MD** (Division of Immunohematology, Ivrea Hospital, Ivrea, Torino); **V. La Milia, MD** (Division of Nephrology and Dialysis, "A. Manzoni" Hospital, Lecco); **C. Marseglia, MD** (Service of Nephrology and Dialysis, "Carlo Poma" Hospital, Mantova); **A. Bettinelli, MD** (Pediatric Division, "S. Leopoldo Mandic" Hospital, Merate, Lecco); **R. Chimenz, MD** (Division of Pediatric Nephrology, "G. Martino" Hospital, Messina); **G. Ardissino, MD, A. Edefonti, MD, C. Fredella, MD, F. Paglialonga, MD** (Division of Pediatric Nephrology, Dialysis and Transplant, "De Marchi" Pediatric Clinic, Milano); **A. Lattuada, BiolSciD., E. Rossi, MD** (Division of Hematology, "L. Sacco" Hospital, Milano); **V. Rossi, MD** (Division of Hematology, "Niguarda Cà Granda" Hospital, Milano); **V. Toschi, MD** (Trasfusioneal Center, "San Carlo Borromeo" Hospital, Milano); **Volpi A, MD** (Division of Nephrology and Dialysis, San Paolo Hospital, Milano); **D. Bonucchi, MD, D. Davoli, MD, M. Leonelli, MD, E. Rubbiani, MD** (Division of Nephrology, Dialysis and Transplant, Policlinico Hospital, Modena); **D. Belotti, BiolSciD., E. Pogliani, MD** (Division of Hematology and Transfusioneal Center, "S. Gerardo" Hospital, Monza, Milano); **G. Masera, MD** (Pediatric Department, "S. Gerardo" Hospital, Monza, Milano); **T. Stellato, MD** (Division of Nephrology, S. Gerardo" Hospital, Monza, Milano); **M.R. Iannuzzi, MD** (Division of Nephrology, "A. Cardarelli" Hospital, Napoli); **A. Ferretti, MD, C. Pecoraro, MD** (Division of Nephrology, Children's Hospital "Santobono", Napoli); **G.B. Capasso, MD, S. Scognamiglio, MD** (Chair of Nephrology, Second University of Napoli, Napoli); **M. Bosa, MD, G.R. Cambrin, MD** (Division of Internal Medicine-Hematology, "S. Luigi" Hospital, Orbassano, Torino); **L. Murer, MD** (Pediatric Division, Policlinico Hospital, Padova); **A. Indovina, MD, R. Marcenò, MD** (Division of Hematology, "V. Cervello" Hospital, Palermo); **L. Amico, MD** (Division of Nephrology and Dialysis, "V. Cervello" Hospital, Palermo); **P. Salis, MD** (ISMETT, Palermo); **U. Maggiore, MD** (Division of Nephrology, Ospedale Maggiore, Parma); **P. Noris, MD, C. Balduini, MD** (Medical Clinic, "San Matteo" Hospital, Pavia); **E. Trabassi, MD** (Division of Nephrology and Dialysis, "San Massimo" Hospital, Penne, Pescara); **G. Agnelli, MD, A. Blass, MD** (Division of Internal Medicine, University of Perugia, Perugia); **I. Capolsini, MD** (Division of Hematology, University of Perugia, Perugia); **P. Poisetti, MD** (Division of Nephrology and Dialysis, Hospital of Piacenza, Piacenza); **R. Caprioli, MD** (Division of Nephrology and Dialysis, "S. Chiara" Hospital, Pisa); **E. Nesti, MD** (Division of Nephrology and Dialysis, "S. Miniato" Hospital, S. Miniato, Pisa); **M. Tuccori, MD** (Division of Farmacology, Azienda Ospedaliera Universitaria Pisana, Pisa); **G. Garozzo, MD** (Trasfusioneal Center, "M.P. Arezzo" Hospital, Ragusa); **E. Bresin, MD, E. Daina, MD, S. Gamba, RN**, (Clinical Research Center for Rare Diseases "Aldo e Cele Daccò", Ranica, Bergamo); **M. Santostefano, MD** (Division of Nephrology and Dialysis, "Santa Maria delle Croci" Hospital, Ravenna); **G. Enia, MD, P. Finocchiario, MD, C. Zoccali, MD** (Division of Nephrology and Dialysis, "Bianchi, Melacrino, Morelli" Hospital, Reggio Calabria); **V. Trapani Lombardo, MD** (Division of Hematology, "Bianchi, Melacrino, Morelli" Hospital, Reggio Calabria); **A. Amendola, MD, L. Dessanti, MD, F. Mandelli, MD, G. Meloni, MD** (Department of Cellular Biotechnology and Hematology, "La Sapienza" University, Roma); **F. Emma, MD, A. Gianviti, MD, S. Rinaldi, MD** (Division of Nephrology and Dialysis, "Bambino Gesù" Pediatric Hospital, Roma); **A. De Feo, MD, M. Ferrannini, MD** (Rome

American Hospital, Roma); **A. Severino, MD** (Division of Hematology, S. Camillo-Forlanini Hospital, Roma); **T. Cicchetti, MD, G. Putorti, MD** (Division of Nephrology and Dialysis, "N. Giannettasio" Hospital, Rossano Calabro, Cosenza); **R. Paolini, MD** (Medical Division, Rovigo Hospital, Rovigo); **A. Pinto, MD** (Division of Nephrology and Dialysis, "S.G. di Dio e Ruggi d'Aragona" Hospital, Salerno); **A. Del Giudice, MD** (Division of Nephrology, "Casa Sollievo delle Sofferenza" Hospital, S. Giovanni Rotondo, Foggia); **P.R. Scalzulli, MD** (Division of Hematology, "Casa Sollievo delle Sofferenza" Hospital, S. Giovanni Rotondo, Foggia); **M. Sanna, MD** (Division of Medical Pathology, Sassari Hospital, Sassari); **A. Amore, MD, G. Conti, MD, R. Coppo, MD, L. Peruzzi, MD** (Division of Nephrology and Dialysis, "Regina Margherita" Pediatric Hospital, Torino); **O. Giacchino, MD, M. Milan, MD** (Service of Immunohematology, "S.G. Bosco" Hospital, Torino); **M. Borca, MD, C. Rollino, MD, A. Vallero, MD** (Service of Nephrology and Dialysis, "S.G. Bosco" Hospital, Torino); **L. Biancone, MD, L. Colla, MD, M. Messina, MD** (Division of Nephrology and Dialysis, "S. G. Battista" Hospital, Torino); **A. Khaled, MD, M. Mazzon, MD, C. Tognoli, MD** (Division of Nephrology, "S. Chiara" Hospital, Trento); **C. Cascone, MD, M. Dugo, MD, S. Mastrosimone, MD** (Division of Nephrology and Dialysis, "S. Maria dei Battisti" Hospital, Treviso); **M. Pennesi, MD** (Division of Pediatric Nephrology, "Burlo Garofolo" Hospital, Trieste); **E. Barbi, MD** (Pediatric Clinic, "Burlo Garofolo" Hospital, Trieste); **G.O. Panzetta, MD** (Division of Nephrology and Dialysis, Hospital of Cattinara, Trieste); **L. Campiotti, MD** (Division of Internal Medicine, "Fondazione Macchi" Hospital, Varese); **O. Amatruda, MD, G. Colussi, MD** (Division of Nephrology, "Fondazione Macchi" Hospital, Varese); **L. Funaro, MD** (Division of Nephrology and Dialysis, "Presidio Ospedaliero", Verbania); **P. Bernich, MD, A. Lupo, MD** (Division of Nephrology and Dialysis, "Borgo Trento" Hospital, Verona); **L. Tavecchia, MD** (Division of Hematology, "Borgo Roma" Hospital, Verona).

Investigators - Abroad

M.G. Caletti, MD, M. Adragua, MD ("Juan P. Garrahan" Hospital de Pediatria, Buenos Aires, Argentina); **P.A. Coccia, MD, J. Ferraris, MD, G. Greloni, MD, R. Groppa, MD, N. Imperiali, MD** (Division of Nephrology, "Hospital Italiano de Buenos Aires", Buenos Aires, Argentina); **P. Henning, MD, J. Nairn, MD** (Renal Unit, Women's and Children's Hospital, North Adelaide, Australia); **R. Wens, MD** (Clinique de Nephrologie-Dialyse, CHU Brugmann, Bruxelles, Belgium); **D. Roussinov, MD** (University Pediatric Hospital, Sofia, Bulgaria); **M. Bitzan, MD** (Pediatric Nephrology, McGill University and Montreal Children's Hospital, Montreal Quebec, Canada); **G. Filler, MD, K. Blyth, RN** (Children's Hospital of Eastern Ontario, Ottawa, Canada); **E. Jančová, MD, R. Ryšavá, MD** (Nephrology Unit, General Faculty Hospital, Praha, Czech Republic); **T. Ring, MD** (Department of Nephrology, Aalborg Hospital, Aalborg, Denmark); **A. Traat, MD** (Tartu University Clinics, Children's Hospital, Tartu, Estonia); **C. Bühner, MD** (Department of Neonatology, Charité Campus Virchow-Klinikum, Berlin, Germany); **D. Müller, MD, U. Querfeld, MD** (Department of Pediatric Nephrology, Charité, Berlin Germany); **B. Hoppe, MD** (University Children's Hospital, Cologne, Germany); **C.V. Schnakenburg, MD** (Department of Pediatrics, University Children's Hospital, Freiburg, Germany); **M.J. Kemper, MD, F. Thaiss, MD** (Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany); **A. Lambert, MD** (Division of Nephrology, University Hospital Mannheim, Germany); **A. Schulze Everding, MD** (Pediatriische Nephrologie, Universitätsklinikum Muenster, Germany); **A. Kattamis, MD, V. Spolou, MD** ("Agia Sophia" Children's Hospital, Athens University, Goudi Athens, Greece); **P. Ziroyannis, MD** (Nephrology Clinic, "G. Gennimatas" General Hospital, Athens, Greece); **P.Y.J. Sim, MD** (Department of Medicine and Geriatrics, Princess Margaret Hospital, Lai Chi Kok, Hong Kong); **M.R. Ardalan, MD** (Nephrology Department, Tabriz Medical University, Tabriz, Iran); **D. Landau, MD** (Division of Pediatric Nephrology,

Soroka Medical Center, Beer-Sheba, Israel); **C. Rinat, MD** (Pediatric Nephrology Unit, Shaare Zedek Medical Center, Jerusalem, Israel); **I. Krause, MD, O. Schiller, MD** (Schneider Children's Medical Center, Petach-Tikvah, Israel); **R. Rahamimov, MD** (Transplantation Department, Beilinson Medical Center, Petach-Tikvah, Israel); **Z. Farfel, MD** (Department of Medicine, Sheba Medical Center, Tel Hashomer, Israel); **E. Matsukuma, MD** (Department of Pediatrics, Nagoya Daini Red Cross Hospital, Nagoya, Japan); **W.J. Wan Ismail, MD** (Pediatric Department, Hospital Selayang, Selangor, Malaysia); **P. Ponce, MD** (Hospital "Garcia de Orta", Almada, Portugal); **J. Barbot, MD, M. Antunes., MD** (Division of Hematology, "Maria Pia" Hospital, Porto, Portugal); **M.S. Faria, MD** (Department of Pediatric Nephrology, "Maria Pia" Hospital, Porto, Portugal); **A.N. Lategann, MD** (AMPATH Laboratories, Arcadia, Republic of South Africa); **M.A. Albalwi, MD, A. Alswaid, MD** (King Abdulaziz Medical City, Riyadh, Saudi Arabia); **S. Al-Saadoun, MD, A. Manlangit, MD** (Pediatric Nephrology, Riyadh Armed Forces Hospital, Saudi Arabia); **R. Bogdanovic, MD** (Department of Nephrology, Institute of Mother and Child Care of Serbia, Belgrade, Serbia); **J.J. Verdu, MD** (Medicina Trasfusional, Hospital General Universitario de Alicante, Spain); **J. Luño Fernandez, MD** (Department of Nephrology, Hospital General Universitario Gregorio Marañón, Madrid, Spain); **H. Cano, MD, C. Martínez, MD** (Centro Regional de Hemodonación, Murcia, Spain); **A. Bock, MD** (Nephrology, Kantonsspital Aarau, Switzerland); **C. Rudin, MD** (University Children's Hospital, Basel, Switzerland); **F. Burkhalter, MD, M. Mayr, MD, M.J. Kim, MD** (Transplantation, Immunology and Nephrology, University Hospital, Basel, Switzerland); **N. Marangon, MD** (Services de Nephrologie et de Transplantation, Hopitaux Universitaires de Geneve, Switzerland); **S. Nef, MD, A. Schenk, MD, G. Spartà, MD** (Nephrology Unit, University Children's Hospital, Zürich, Switzerland); **B. Bergamin, MD, T. Fehr, MD** (University Hospital, Zürich, Switzerland); **E. Yilmaz Keskin, MD** (Pediatric Hematology Unit, Gazi University, Ankara, Turkey); **A. Sharma, MD** (Royal Liverpool and Broadgreen University Hospitals, Liverpool, United Kingdom); **D. Bockenbauer, MD** (Great Ormond Street Hospital, London, United Kingdom); **G.B. Haycock, MD** (Pediatric Renal Unit, Guy's Hospital, London, United Kingdom); **M. Ognjanovic, MD** (Royal Victoria Infirmary, Newcastle upon Tyne, United Kingdom); **H. Gallagher, MD** (South West Thames Renal and Transplantation Unit, St. Helier Hospital, Surrey, United Kingdom); **A. Katz, MD** (Pediatric Nephrology, Children's Hospital, Birmingham, AL, USA); **J. Mittleman, MD** (Cedars-Sinai Medical Center, Los Angeles, CA, USA); **S. Godfrey, MD** (Kaiser Hospital, San Diego, CA, USA); **V.R. Dharnidharka, MD** (University of Florida, Division of Pediatric Nephrology, Gainesville, FL, USA); **E. Chang, MD** (St Luke's Mountain States Tumor Institute, Pediatric Hematology, Boise, ID, USA); **J.C. Lane, MD, C.B. Langman, MD** (Division of Kidney Diseases, Children's Memorial Hospital, Chicago, IL, USA); **V. Kimonis, MD** (Department of Pediatrics, SIU School of Medicine, Springfield, IL, USA); **L. Milner, MD** (Division of Nephrology, Floating Hospital for Children, Boston, MA, USA); **R. Montgomery, MD** (Johns Hopkins Hospital, Baltimore, MD, USA); **M. Hand, MD** (Pediatric Nephrology, The Barbara Bush Children's Hospital at Maine Medical Center, ME, USA); **C.E. Kashtan** (Pediatrics, University of Minnesota Medical School, Minneapolis, USA); **L. Najera, MD, J. Steinke, MD** (Pediatric Nephrology, Fairview University Medical Center, Minneapolis, USA); **D. Milliner, MD** (Mayo Clinic, Rochester, MN, USA); **B. Warady, MD** (Children's Mercy Hospital, Kansas City, MO, USA); **M. Ferris, MD, D. Bunch, PhD** (Division of Nephrology and Hypertension, University of North Carolina, Chapel Hill, NC, USA); **P. Lane, MD, L. Wrenshall, MD** (Nebraska Medical Center, Omaha, NE, USA); **K. Lieberman, MD** (Pediatric Nephrology, Hackensack University Medical Center, Hackensack, NJ, USA); **R. Wallerstein, MD** (Genetics Department, Hackensack University Medical Center, Hackensack, NJ, USA); **R. Tapia, MD** (Division of Pediatric Nephrology, The Children's Hospital at Albany Medical Center, Albany, NY, USA); **J. Listman, MD** (SUNY Upstate Medical University, Syracuse, NY, USA); **S.B. Conley, MD**

(Department of Nephrology, St. Christopher's Hospital for Children, Philadelphia, USA); **A.K. Feng, MD** (Pediatric Critical Care, Hasbro Children's Hospital, Providence, RI, USA); **M.G. Seikaly, MD** (Pediatric Nephrology, Children's Medical Center, Dallas, Texas, USA); **R. Raafat, MD** (Pediatric Nephrology, University of Texas Health Science Center, San Antonio, Texas, USA); **J.C. Barrett, MD** (Division of Hematology-Oncology, Virginia Commonwealth University, Richmond, VA, USA); **V. Chadha, MD** (VCU Medical Center, Pediatric Nephrology, Richmond, VA, USA); **J. Felgenhauer, MD** (Pediatric Oncology Center, Sacred Heart Children's Hospital, Spokane, WA, USA); **J. Gitomer, MD** (Department of Nephrology, Marshfield Clinic, Marshfield, WI, USA).

Laboratory Analysis

F. Gaspari, ChemD. (Clinical Research Center for Rare Diseases "Aldo e Cele Daccò", Ranica, Bergamo); **C. Ottomano, MD, A. Vernocchi, MD** (Division of Laboratory Analysis, "Ospedali Riuniti" Azienda Ospedaliera Bergamo).

Biochemical Studies

P. Bettinaglio, BioSciD, J. Caprioli, BioSciD, F. Castelletti, BiotechD, D. Cugini, BioSciD, C. Mele, BioSciD, G. Pianetti, Chemist (Clinical Research Center for Rare Diseases "Aldo e Cele Daccò", Ranica, Bergamo); **M. Galbusera, BioSciD, S. Gastoldi, Chemist, D. Macconi, BioSciD** ("M. Negri" Institute for Pharmacological Research, Bergamo, Italy); **P.F. Zipfel, MD** (Hans Knoell Institute for Natural Products Research, Jena, Germany).

Statistical Analysis

A. Perna, StatSciD (Clinical Research Center for Rare Diseases "Aldo e Cele Daccò", Ranica, Bergamo).